

# Note On Endomorphism Algebras Of Separable Monoidal Functors.

Brian J. DAY and Craig A. PASTRO  
April 26, 2009.

*Abstract: We recall the Tannaka construction in [2] for certain types of split monoidal functor into  $\mathbf{Vect}_k$ , and remove the “compactness” restriction on the domain.*

In  $\mathbf{Set}$  or  $\mathbf{Vect}_k$ , a “VN-core” is an algebra  $(A, \mu, \eta)$  and a coalgebra  $(A, \delta, \epsilon)$  satisfying the axiom

$$\delta\mu = (\mu \otimes \mu)(1 \otimes c \otimes 1)(\delta \otimes \delta),$$

where  $c$  is the symmetry for  $\otimes$ , with a map  $S : A \rightarrow A$  such that

$$\mu_3(1 \otimes S \otimes 1)\delta_3 = 1 : A \rightarrow A.$$

The core is called unital if it satisfies the stronger axiom

$$1 \otimes \eta = (1 \otimes \mu)(1 \otimes S \otimes 1)\delta_3.$$

In  $\mathbf{Set}$ , unital VN-cores are precisely groups, while in  $\mathbf{Vect}_k$  they contain the Hopf  $k$ -algebras.

The following theory is  $k$ -linear for  $k$  a field of characteristic 0, and we shall assume some familiarity with [2] Theorem 3.1 which we now generalize. The aim of this procedure is to remove the construction of the unital VN-core  $\mathbf{End}^\vee U$  in [2] from the “compact” setting by postulating the existence of an abstract duality functor

$$(-)^* : \mathcal{A}^{\text{op}} \rightarrow \mathcal{A}$$

on the generating category  $\mathcal{A} \subset \mathcal{C}$  with an associated natural isomorphism

$$u : U(A^*) \cong U(A)^*$$

and a transformation  $e : A^* \otimes A \rightarrow I$  satisfying the two axioms

$(e, r, r_0)$ :

$$\begin{array}{ccc}
 k & \xrightarrow{\quad r_0 \quad} & UI \\
 \uparrow e & & \uparrow Ue \\
 U(A)^* \otimes UA & & U(A^* \otimes A) \\
 & \swarrow u \otimes 1 \quad \searrow r & \\
 & U(A^*) \otimes UA & 
 \end{array}$$

commutes, and

$(e, i, i_0)$ :

$$\begin{array}{ccc}
 UI & \xrightarrow{i_0} & k \\
 \uparrow Ue & & \uparrow e \\
 U(A^* \otimes A) & & U(A)^* \otimes UA \\
 & \searrow i \quad \nearrow u \otimes 1 & \\
 & U(A^*) \otimes UA &
 \end{array}$$

commutes.

We emphasize here that it is not necessary for  $A^*$  to be the actual  $\otimes$ -dual of  $A$  in the monoidal category  $\mathcal{C}$ . For example,  $\mathcal{C}$  could be a  $\star$ -autonomous monoidal category [1], or just a monoidal closed category [3], and so forth.

The two axioms above now replace the cumbersome “ $U$ -trace” condition in [2] § 3. The region labeled (3) in [2] § 3 now becomes:

$$\begin{array}{ccccc}
 & & (U(A)^* \otimes UA)^* \otimes U(A)^* \otimes UA & & \\
 & \nearrow e^* \otimes 1 \otimes 1 & & \searrow (u \otimes 1)^* \otimes (u^{-1} \otimes 1) & \\
 k^* \otimes U(A)^* \otimes UA & & & & (U(A^*) \otimes UA)^* \otimes U(A^*) \otimes UA \\
 \uparrow 1 \otimes c & \searrow 1 \otimes u^{-1} \otimes 1 & & \downarrow i^* \otimes r & \\
 k^* \otimes UA \otimes U(A)^* & & k^* \otimes U(A^*) \otimes UA & & \\
 \uparrow 1 \otimes e & \searrow 1 \otimes e & \downarrow i_0^* \otimes r & \nearrow U(e)^* \otimes 1 & \\
 k^* \otimes k & & U(I)^* \otimes U(A^* \otimes A) & & U(A^* \otimes A)^* \otimes U(A^* \otimes A) \\
 \uparrow 1 & \searrow i_0^* \otimes r_0 & \downarrow 1 \otimes U(e) & \nearrow nat. & \downarrow copr_{B=A^* \otimes A} \\
 k^* \otimes k & \xrightarrow{i_0^* \otimes r_0} & U(I)^* \otimes UI & \xrightarrow{copr_{B=I}} & \int^B U(B)^* \otimes UB
 \end{array}$$

which eventually commutes by commutativity of

$$\begin{array}{ccc}
 UA \otimes U(A)^* & \xrightarrow{c} & U(A)^* \otimes UA \\
 \uparrow n & (trace) & \downarrow e \\
 k & \xrightarrow{(\dim UA).1} & k
 \end{array}$$

in  $\mathbf{Vect}_k$ .

Thus we essentially replace the map  $S$  in [2] Theorem 3.1 by the map defined on  $\mathbf{End}^\vee U$  by the new family:

$$S_A = (\dim UA)^{-1} \cdot \sigma_A$$

(i.e. remove the factor  $(\dim UI)$  and assume  $\dim UA \neq 0$  for all  $A \in \mathcal{A}$ ). The new structure  $(\mathbf{End}^\vee U, \mu, \eta, \delta, \epsilon, S)$  is then a unital VN-core in  $\mathbf{Vect}_k$ . Note that we no longer require  $\mathcal{C}$  and  $U$  to be braided structures in this context.

## References.

- [1] M. Barr, “ $\star$ -Autonomous Categories”,  
Lecture Notes in Mathematics, (Springer) 752 (1979).
- [2] B. J. Day and C. A. Pastro, “On Endomorphism Algebras of Separable Monoidal Functors”,  
Theory Appl. Categories, Vol. 22, No. 4 (2009), Pg. 77-96. (Electronic).
- [3] S. Eilenberg and G. M. Kelly, “Closed Categories”,  
Proc. Conf. on Categorical Algebra, La Jolla 1965, Springer-Verlag (1966) Pg. 421-562.

Mathematics Dept., Faculty of Science, Macquarie University, NSW 2109, Australia.

Any replies are welcome through Tom Booker (thomas.booker@students.mq.edu.au),  
who kindly typed the manuscript.